

**Joint Replacement, Sports Medicine,
Fracture Care & General Orthopaedics**

Robert F. Jones, MD
Marisa A. Schmitt, MD
Alison Bird, MHP, PA-C



**Phone (978) 534-6333
www.longviewortho.com**

100 Hospital Road, Suite 3C
Leominster, MA 01453
Fax (978) 840-0966

SHOULDER SURGERY - Postoperative Instructions

- Rotator Cuff Repair** **Biceps tenotomy/tenodesis** **Labral debridement**
 Subacromial decompression **Excision distal clavicle** **Loose body removal**

Follow-up appointment: Our office will call you the next business day after surgery to schedule a postoperative visit (usually 10 days postop). At that time you will be given the opportunity to ask questions about your surgery or discuss any difficulties you may have. Photographs from arthroscopic procedures may be reviewed at the follow-up visit.

About your dressing: Your shoulder is covered with a sterile dressing following the procedure, and this should be kept clean and dry for 48 hours. You may shower or bathe, but be sure that the dressing is covered and kept out of the flow of water. Drainage is very common following shoulder surgery so you may need to change the dressing if it becomes soiled or wet. Remove the dressing after two days and cover the wound with gauze or a band aid. The stitches used to close the wounds are made of nylon and will be removed when you return for your follow-up visit.

About pain and swelling: Some discomfort is to be expected following this surgery. Your surgeon uses a local anesthesia surrounding the incision that will provide pain relief for 6-10 hours following surgery. Following shoulder surgery, patients often find that sleeping is more comfortable in a propped up, semi-sitting position (i.e. sleeping in a recliner or in bed with multiple pillows).

For pain, use the narcotic medication that your doctor ordered. Please do not combine narcotics with alcoholic beverages.

Nonsteroidal anti-inflammatories (like Motrin, Advil or Aleve) may be used with or instead of the narcotic medication.

- Use ice (in a plastic bag wrapped in a dry towel), 20 minutes out of an hour, repeating as needed
 Use Polar Care device – instructions included

Postoperative activities:

Use the sling/immobilizer to help minimize pain following surgery. You may remove the sling for gentle range of motion exercises to the shoulder as tolerated. Discontinue the sling when comfort allows.

Leave the immobilizer in place except to shower and change clothing. **Do not actively move your shoulder,** but it can be moved gently with gravity or assistance from the other arm to help with dressing/showering. It is recommended to remove the sling at least 3 times each day for Range of Motion to the fingers/wrist/elbow to prevent those joints from getting stiff.

Use bolster pillow at all times/if comfortable

Return to driving/work:

Return to work will depend on your work duties- this should be discussed with your surgeon.

Return to driving:

- Allowed when no **longer taking narcotics and can comfortably reach the steering wheel**
 Not allowed until cleared by your surgeon

Warning signs: If you have fevers or chills, the pain is not manageable, there is excessive bloody drainage or signs of infection, or if you have any other questions, **please call our office at #978-534-6333.**